

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Stellar Populations in the Magellanic Clouds

Nino Panagia¹, Guido De Marchi², Martino Romaniello³

(1) *Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA; also, INAF-HQ, Via del Parco Mellini 84, 00136 Rome, Italy; and Supernova Ltd., OYV #131, Northsound Road, Virgin Gorda, British Virgin Islands*

(2) *ESA/ESTEC, Keplerlaan 1, 2200 AG Noordwijk, The Netherlands*

(3) *ESO, Karl-Scharzschild-Strasse 2, 85748 Garching-bei-München, Germany*

Abstract. We present the first results of our study of stellar populations in the Large and Small Magellanic Clouds based on multiband WFPC2 observations of “random” fields taken as part of the “pure parallel” program carried out by the Space Telescope Science Institute as a service to the community.

We have started a study of the stellar populations in the Large and Small Magellanic Clouds using data (U, B, V, I and H α imaging) collected as part of the WFPC2 Pure Parallel Program (Wadadekar et al. 2006). We have considered a number of fields in both Magellanic Clouds which comprise stellar populations with ages ranging from a few Myr to a few Gyr and in which pre-main-sequence objects are clearly identified. In addition to the properties of these populations, we have studied the characteristics of the dust present in these regions.

Using the properties of the red giants of the so-called “red clump” (RC) in the various colour-magnitude diagrams, we have determined the extinction law in each field (Panagia & De Marchi 2005). We find that not only the amount of extinction but also its wavelength dependence varies considerably from site to site, indicating that the physical properties of the absorbing dust are not uniform across the Clouds (see Figure 1). This analysis also allows us to determine the relative three-dimensional distribution of the different populations as well as their location with respect to the absorbing dust clouds.

Correcting for the appropriate extinction for each field *individually*, and following the procedures developed by Panagia et al. (2000), Romaniello et al. (2002), and Romaniello, Robberto & Panagia (2004), we determine the physical parameters, namely effective temperature and luminosity, of all stars present in the field (De Marchi et al. 2005). Table 1 illustrates some of the properties of the stellar populations for the fields we have analyzed so far.

We find that, among young stars, those less massive than 2 solar masses are spatially more spread over each field than massive B type stars, whereas old stars are quite uniformly distributed. The data also reveal that young stars, which appear to be spatially associated with the nebular emission, are physically confined within a small range of distances, whereas old stars are uniformly distributed at all depths along the line of sight. An important corollary of our investigation is that unaccounted patchy absorption or a variable extinction law

are likely to contribute significantly to the present discrepancies between various distance indicators for the Magellanic Clouds.

References

- De Marchi, G., Panagia, N., & Kozhurina-Platais, V., & Romaniello, M. 2005, BAAS, 37, 1368 ; and in preparation.
 Panagia, N., Romaniello, M., Scuderi, S., & Kirshner, R.P. 2000, ApJ, 539, 197.
 Panagia, N., & De Marchi, G. 2005, BAAS, 37, 1368; and in preparation.
 Romaniello, M., Panagia, N., Scuderi, S., & Kirshner, R.P. 2002, AJ, 123, 215.
 Romaniello, M., Robberto, M., & Panagia, N. 2004, ApJ, 608, 220.
 Wadadekar, Y., et al. 2006, PASP, 118, 450.




Figure 1. Reddening curves for several fields in the LMC and SMC compared with the standard Milky Way extinction law (dotted lines).

Table 1. Properties of Stellar Populations in several LMC and SMC Fields

Field	E(V-I) range	$\langle A_V \rangle$	V_{min} brightest young star	M_{max} highest mass star [$M_{\odot}$]	T_{min} youngest blue star [Myr]	T_{RC} min age RC [Gyr]	Z_{RC} metallicity range RC [$Z_{\odot}$]
LMC1	0.00-0.06	0.14	15.5	5	100	1.4	0.2-0.4
LMC2	0.07-0.20	0.50	16.1	6	70	1.0	0.2-0.6
LMC7	0.30-0.70	1.95	13.8	36	4	0.5	0.2-0.6
SN87A	0.20-0.50	1.23	13.3	14	16	0.5	0.2-0.4
SMC1	0.04-0.08	0.21	15.8	9	30	1.4	<0.2
SMC2	0.08-0.16	0.42	16.5	7	50	1.4	<0.2